



The challenges in classifying rheumatoid arthritis-associated interstitial lung disease

Hanna Nurmi^{1,2}, Riitta Kaarteenaho³

¹Division of Respiratory Medicine, Institute of Clinical Medicine, School of Medicine, Faculty of Health Sciences, University of Eastern Finland, Kuopio, Finland; ²Center of Medicine and Clinical Research, Division of Respiratory Medicine, Kuopio University Hospital, Kuopio, Finland; ³Research Unit of Internal Medicine, University of Oulu & Medical Research Center (MRC) Oulu, Oulu University Hospital, Oulu, Finland

Correspondence to: Prof. Riitta Kaarteenaho, MD, PhD. Research Unit of Internal Medicine, University of Oulu & Medical Research Center (MRC) Oulu, Oulu University Hospital, Kajaanintie 50, 90220 Oulu, Finland. Email: Riitta.Kaarteenaho@oulu.fi.

Provenance and Peer Review: This article was commissioned by the editorial office, *Journal of Thoracic Disease*. The article did not undergo external peer review.

Comment on: Yamakawa H, Sato S, Tsumiyama E, *et al.* Predictive factors of mortality in rheumatoid arthritis-associated interstitial lung disease analysed by modified HRCT classification of idiopathic pulmonary fibrosis according to the 2018 ATS/ERS/JRS/ALAT criteria. *J Thorac Dis* 2019;11:5247-57.

Submitted Feb 11, 2020. Accepted for publication Mar 06, 2020.

doi: 10.21037/jtd.2020.03.96

View this article at: <http://dx.doi.org/10.21037/jtd.2020.03.96>

Interstitial lung diseases (ILD) constitute a large group of disorders with variable prognoses and treatment options (1). The current international guidelines on idiopathic interstitial pneumonias (IIP) and idiopathic pulmonary fibrosis (IPF), the most common type of IIP, have focused on the classification of subtypes based on high-resolution computed tomography (HRCT) patterns (2,3). At present, no similar classification guidelines to those for IPF and IIP have been issued for connective tissue disease (CTD)-related, such as rheumatoid arthritis-associated ILD (RA-ILD). Instead, these classification guidelines of IIP and IPF have often been adopted in clinical practice. Due to the risks associated with surgical lung biopsies, most patients are diagnosed and classified into the different subtypes based only on HRCT (4,5). A very recent review has nicely summarized studies reporting RA-ILD HRCT patterns and mortality (6).

Predicting the survival of the individual patient with ILD is extremely challenging, and only a few studies have explored this issue in RA-ILD (7). Thus, clinicians have very few tools to assist them when making a prognosis estimation in patients suffering from RA-ILD. Previously, factors reflecting the RA severity have been proposed to associate with worse survival, including increased erythrocyte sedimentation rate and high visual analogue

pain scale, etc. (7). Male sex, age and both baseline lung function test results and the longitudinal change of their values have also associated with mortality (7-9).

In addition, some previous studies have investigated the prognostic role of HRCT and revealed that the extent of the radiologic ILD changes was related with mortality (10,11), and that the usual interstitial pneumonia pattern (UIP) in HRCT associated with worse survival than the other subtypes (12-14), although this has not been detected by all investigators (15). Some findings from our recent study revealed that the extents of reticulation, traction bronchiectasis and architectural distortion were associated with decreased survival, whereas the presence or extent of honeycombing was not beneficial in predicting survival, even though it correlated with hospitalizations due to respiratory reasons (16). Our finding contrasts with the study of Kim *et al.* in which the extents of honeycombing and traction bronchiectasis were independent predictors of worse outcome (12).

Overall, the studies regarding to the predictive value of HRCT in RA-ILD are sparse and partly controversial. Therefore, we read the article of Hideaki Yamakawa and co-authors with much interest and thank the authors for investigating this important topic (17).

They have analyzed the medical records of 96 patients

with RA-ILD using combined modified IPF (ATS/ERS 2018) and IIP (ATS/ERS 2013) guidelines for the classification of the cases, with the aim of identifying factors that would predict mortality by examining HRCT patterns. RA diagnoses were confirmed by a rheumatologist and a positive anticyclic citrullinated peptide (anti-CCP) value. In the study protocol, each subject's radiological findings were reviewed by two expert pulmonologists, but not by radiologists. Disagreements between the two pulmonologists were resolved by discussion and the interrater agreement between them was shown to be good (kappa value 0.75).

The cases were categorized based on HRCT into seven subgroups, namely definite UIP, probable UIP, indeterminate for UIP, non-specific interstitial pneumonia (NSIP), organizing pneumonia (OP), NSIP + OP and unclassifiable patterns. The authors concluded that indeterminate for UIP pattern (30%) was the most common type. One may speculate whether this conclusion was clinically meaningful, since if one combined the definite UIP (21%) and the probable UIP patterns (20%) then this accounted for more than 40% of the cases.

Different subtypes did not correlate positively with survival, which was in contrast to the results of some previous studies (8,12,14), whereas the existence of honeycombing did show a correlation. One may wonder, however, if the analyses had been performed differently, e.g., by comparing definite UIP cases with all other cases, or definite and probable UIP cases together with all others, then this might have produced somewhat different results. Indeed, in the rather recent longitudinal study of Yunt *et al.* the combined group of definite and possible UIP patterns had a significantly worse survival than the NSIP patients (18). Moreover, as the disease progressed, almost every second patient originally displaying a probable UIP pattern developed a definite UIP pattern (19). In addition, Yamakawa's group has recently published another study focusing on honeycombing in RA-ILD, revealing that honeycomb formation occurred in 40% of those RA-ILD patients in whom it was not present at baseline HRCT (20). Analyzing the longitudinal change or the extent of the radiological changes would have strengthened the present study even more. Furthermore, it remained unclear how the existence of honeycombing was detected—by reviewing the radiological reports or by re-analyzing HRCTs?

The question of distribution of the ILD changes is important but difficult to resolve. In idiopathic ILD forms, the distribution in the definite UIP pattern is typically subpleural and basal predominant, but often heterogeneous

(2,3). In RA-ILD, the distribution of disease may not display a basal predominance but reticulation and honeycombing can be concentrated in the middle or upper zones of the lungs (21). In the recently published study of Jacob *et al.*, RA-ILD patients with a definite UIP pattern regardless of whether the distribution was IPF-like or not, demonstrated a similar outcome (22). In the study on which we are commenting here, the NSIP/UIP category was selected if the distribution was central/diffuse even when reticulation with traction bronchiectasis and honeycombing were present and ground-glass opacities were inconspicuous (17). Perhaps these patients could have been categorized as definite UIP? Honeycombing was present only in the definite UIP and in the NSIP/UIP subgroups. Some patients in the latter subgroup could just as correctly have been categorized as definite UIP, which might have changed the results.

We congratulate the authors for the clever study protocol and for compilation of the modified classification; however this would have been even more valuable if the HRCT scans had been re-analyzed by thoracic radiologists. This fact was also stated by the authors themselves who have mentioned in the discussion as the limitation of their study as follows: “*Expert thoracic radiologists should analyze HRCT findings, not pulmonologists*”. Several previous studies have shown that the repeatability between radiologists on HRCT patterns analyses in IPF is low (23-25). Thus, one may assume that the repeatability between radiologists in a modification involving two distinct guidelines, as done in the present study and examining several IIP patterns would be even lower than simply analyzing IPF.

The authors have, however, presented a suitable combination of the two present classifications of IIP and IPF. Hopefully this will increase the interest of experts in this field in developing its own specific radiological classification for RA-ILD, which would be beneficial not only for research purposes but would also be of assistance in clinical practice.

Acknowledgments

The authors wish to thank Ewen MacDonald for providing assistance with the language.

Funding: None.

Footnote

Conflicts of Interest: Both authors have completed the

ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/jtd.2020.03.96>). HN has been a lecturer/organizer in educational gatherings sponsored by medical companies (Boehringer-Ingelheim, Roche), received congress travel costs (Boehringer-Ingelheim, Roche, Sanofi Genzyme) outside the submitted work and received grants for scientific work from the Kuopio region Respiratory Foundation and the North Savo Regional Fund of the Finnish Cultural Foundation. RK has received consultant fees from GlaxoSmithKline and Boehringer-Ingelheim, lecture fees from Roche and Boehringer-Ingelheim, and a congress travel cost from Orion outside the submitted work, as well as grants for the study group from Foundation of the Finnish Anti-Tuberculosis Association, the Research Foundation of the Pulmonary Diseases, Jalmari and Rauha Ahokas Foundation and the Research Foundation of North Finland.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any parts of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. American Thoracic Society, European Respiratory Society. American Thoracic Society/European Respiratory Society International Multidisciplinary Consensus Classification of the Idiopathic Interstitial Pneumonias. This joint statement of the American Thoracic Society (ATS), and the European Respiratory Society (ERS) was adopted by the ATS board of directors, June 2001 and by the ERS Executive Committee, June 2001. *Am J Respir Crit Care Med* 2002;165:277-304.
2. Raghu G, Remy-Jardin M, Myers JL, et al. Diagnosis of Idiopathic Pulmonary Fibrosis. An Official ATS/ERS/JRS/ALAT Clinical Practice Guideline. *Am J Respir Crit Care Med* 2018;198:e44-68.
3. Travis WD, Costabel U, Hansell DM, et al. An official American Thoracic Society/European Respiratory Society statement: Update of the international multidisciplinary classification of the idiopathic interstitial pneumonias. *Am J Respir Crit Care Med* 2013;188:733-48.
4. Kaarteenaho R. The current position of surgical lung biopsy in the diagnosis of idiopathic pulmonary fibrosis. *Respir Res* 2013;14:43.
5. Raghu G, Remy-Jardin M, Myers J, et al. The 2018 Diagnosis of Idiopathic Pulmonary Fibrosis Guidelines: Surgical Lung Biopsy for Radiological Pattern of Probable Usual Interstitial Pneumonia Is Not Mandatory. *Am J Respir Crit Care Med* 2019;200:1089-92.
6. Bendstrup E, Møller J, Kronborg-White S, et al. Interstitial Lung Disease in Rheumatoid Arthritis Remains a Challenge for Clinicians. *J Clin Med* 2019. doi: 10.3390/jcm8122038.
7. Assayag D, Lubin M, Lee JS, et al. Predictors of mortality in rheumatoid arthritis-related interstitial lung disease. *Respirology* 2014;19:493-500.
8. Solomon JJ, Chung JH, Cosgrove GP, et al. Predictors of mortality in rheumatoid arthritis-associated interstitial lung disease. *Eur Respir J* 2016;47:588-96.
9. Song JW, Lee HK, Lee E, et al. Clinical course and outcome of rheumatoid arthritis-related usual interstitial pneumonia. *Sarcoidosis Vasc Diffuse Lung Dis* 2013;30:103-12.
10. Lee H, Choe J, Kim S, et al. Important Prognostic Factor in Rheumatoid Arthritis Patients with Interstitial Lung Disease Is Not Usual Interstitial Pneumonia Pattern but Interstitial Lung Disease Extent On Chest High-Resolution Computed Tomography. *Arthritis Rheum* 2012;64:S528.
11. Sathi N, Urwin T, Desmond S, et al. Patients with limited rheumatoid arthritis-related interstitial lung disease have a better prognosis than those with extensive disease. *Rheumatology* 2011;50:620.
12. Kim EJ, Elicker BM, Maldonado F, et al. Usual interstitial pneumonia in rheumatoid arthritis-associated interstitial lung disease. *Eur Respir J* 2010;35:1322-8.
13. Akira M, Sakatani M, Hara H. Thin-section CT findings in rheumatoid arthritis-associated lung disease: CT patterns and their courses. *J Comput Assist Tomogr* 1999;23:941-8.
14. Tsuchiya Y, Takayanagi N, Sugiura H, et al. Lung diseases directly associated with rheumatoid arthritis and their relationship to outcome. *Eur Respir J* 2011;37:1411-7.
15. Nurmi HM, Purokivi MK, Kärkkäinen MS, et al. Variable

- course of disease of rheumatoid arthritis-associated usual interstitial pneumonia compared to other subtypes. *BMC Pulm Med* 2016;16:107.
16. Nurmi HM, Kettunen H, Suoranta S, et al. Several high-resolution computed tomography findings associate with survival and clinical features in rheumatoid arthritis-associated interstitial lung disease. *Respir Med* 2018;134:24-30.
 17. Yamakawa H, Sato S, Tsumiyama E, et al. Predictive factors of mortality in rheumatoid arthritis-associated interstitial lung disease analysed by modified HRCT classification of idiopathic pulmonary fibrosis according to the 2018 ATS/ERS/JRS/ALAT criteria. *J Thorac Dis* 2019;11:5247-57.
 18. Yunt ZX, Chung JH, Hobbs S, et al. High resolution computed tomography pattern of usual interstitial pneumonia in rheumatoid arthritis-associated interstitial lung disease: Relationship to survival. *Respir Med* 2017;126:100-4.
 19. Salvatore M, Singh A, Yip R, et al. Progression of probable UIP and UIP on HRCT. *Clin Imaging* 2019;58:140-4.
 20. Yamakawa H, Sato S, Nishizawa T, et al. Impact of radiological honeycombing in rheumatoid arthritis-associated interstitial lung disease. *BMC Pulm Med* 2020;20:25.
 21. Rajasekaran BA, Showlin D, Lord P, et al. Interstitial lung disease in patients with rheumatoid arthritis: A comparison with cryptogenic fibrosing alveolitis. *Rheumatology* 2001;40:1022-5.
 22. Jacob J, Hirani N, van Moorsel, et al. Predicting outcomes in rheumatoid arthritis related interstitial lung disease. *Eur Respir J* 2019. doi: 10.1183/13993003.00869-2018.
 23. Lynch DA, Godwin JD, Safrin S, et al. High-resolution computed tomography in idiopathic pulmonary fibrosis: Diagnosis and prognosis. *Am J Respir Crit Care Med* 2005;172:488-93.
 24. Gruden JF, Panse PM, Gotway MB, et al. Diagnosis of usual interstitial pneumonia in the absence of honeycombing: Evaluation of specific CT criteria with clinical follow-up in 38 patients. *AJR Am J Roentgenol* 2016;206:472-80.
 25. Watadani T, Sakai F, Johkoh T, et al. Interobserver variability in the CT assessment of honeycombing in the lungs. *Radiology* 2013;266:936-44.

Cite this article as: Nurmi H, Kaarteenaho R. The challenges in classifying rheumatoid arthritis-associated interstitial lung disease. *J Thorac Dis* 2020;12(6):3000-3003. doi: 10.21037/jtd.2020.03.96